

# Comparison of Atmospheric Parameters

## Derived from *In-Situ* and Hyperspectral Remote Sensing Data of Beautiful Bavarian Lakes

S. Riedel<sup>1,2</sup>, P. Gege<sup>2</sup>, M. Schneider<sup>2</sup>, B. Pflug<sup>2</sup>, N. Oppelt<sup>1</sup>

### Test Sites and Goal

**Test area:** lake Starnberg and Klostersee (near Munich)

**Goal:** compare derived parameters (Aerosol optical thickness, Angstrom exponent, water vapor column and ozone column) from *in-situ* data collected during a measurement campaign with spectra and parameters derived from HySpex (hyperspectral airborne imager) and Sentinel-2A images.

**Field instruments:** Microtops sunphotometer, Ibsen field spectrometer, RAMSES (TriOS) system

**Imaging sensors:** HySpex (Klostersee), Sentinel-2A (Lake Starnberg)




Fig. 1: **Left:** section of the used Sentinel-2A scene containing Lake Starnberg. **Right:** section of the used HySpex scene containing the Klostersee. Test areas are marked with a red square.

### Conclusion

- Aerosol parameters are generally in good agreement
- High differences in H<sub>2</sub>O and O<sub>3</sub> absorption between *in-situ* and hyperspectral data
- Validation of atmospheric parameters derived from multi-/hyperspectral remote sensing data with *in-situ* measurements is very valuable
- Final interpretation of the observed differences requires further investigations



Fig. 2: The *in-situ* measurement team while preparing the boat at the Lake Starnberg test site. The RAMSES system is suspended from the white crane-like arm.

### Deriving atmospheric parameters from in-situ and remote sensing data




Fig. 3: Aerosol optical thickness (AOT) values at different wavelengths and times derived from Microtops (sunphotometer) measurements. For comparison two Cimel sunphotometer measurements (Aeronet station at Hohenpeissenberg) are included into the plot.




Fig. 4: Downwelling irradiance ( $E_D$ ) measurements for validation of the Lake Starnberg Sentinel-2A data. **Red** curve: data from the RAMSES irradiance sensor (resolution<sub>FWHM</sub> = 12 nm; 3.3 nm sampling distance). **Blue** curve: Downwelling irradiance on a spectralon panel measured with an Ibsen spectrometer (resolution<sub>FWHM</sub> = 1.6 nm; 0.5 nm sampling distance).




Fig. 5: **(a)** Water vapor map of Lake Starnberg (left) derived from Sentinel-2A data and Klostersee (right) derived from HySpex data. Over the water areas water vapor is set constant. **(b)** Same as (a) for Aerosol optical thickness at 550 nm.

**(a) and (b):** average values for comparison have been derived over land surface close to the test areas.

Custom analytical evaluation and Aeronet<sup>(4)</sup> website

AOT 550 nm  
Angstrom exp.  
Water vapor  
Ozone

WASI<sup>(1)</sup> fitting

Water vapor  
Ozone

ATCOR<sup>(3)</sup> and  
Sen2Cor<sup>(2)</sup>

AOT 550 nm  
Water vapor

### Comparison of atmospheric parameters

Lake Starnberg: comparison of atmospheric parameters Sentinel-2A overflight 13.08.2015, 10:16 UTC					
	Microtops (10:17)	Ibsen $E_d$ (10:08)	Ramses $E_d$ (9:54)	Sentinel-2A (10:16)	Aeronet (10:22)
AOT 550 nm	0.151 (interpolated)	-	-	0.205±0.003	0.168 (interpolated)
Angstrom exponent	1.89	-	-	-	1.94 (440-675nm)
Water vapor column	0.93	1.03±0.01	1.02	1.52±0.12	1.44
Ozone column	0.324±0.002	0.459±0.07	0.398	-	-

Table 1: **Left:** parameter comparison for the Lake Starnberg test site. **Right:** parameter comparison for the Klostersee test site. The colored values denote the credibility of the value judged by the authors. **Red:** likely erroneous. **Orange:** maybe erroneous. **Green:** probably close to reality.

Klostersee: comparison of atmospheric parameters Hyperspectral overflight 12.08.2015, 8:57 and 9:04 UTC				
	Microtops (8:55/9:09)	Ramses $E_d$ (9:46)	Hyperspectral (8:57/9:04)	Aeronet (8:52/9:07)
AOT 550 nm	0.212/0.205 (interpolated)	-	0.197±0.003/ 0.192±0.003	0.260/0.253 (interpolated)
Angstrom exponent	1.83±0.01/ 1.75	-	-	1.84/1.86 (440-675nm)
Water vapor column	0.96±0.02/ 0.99	0.93±0.01	2.06±0.07/ 2.02±0.07	1.69
Ozone column	0.323±0.002/ 0.326	0.387±0.002	-	-

- Generally good agreement for aerosol parameters
- Not very good agreement for H<sub>2</sub>O and O<sub>3</sub> absorption
- Values derived from hyperspectral data always higher than *in-situ* values

- Errors are statistical, do not reflect systematic instrument errors
- Values without errors from measurements, upon which no statistics can be applied
- Aeronet distance to Lake Starnberg: 23 km; to Klostersee: 78 km

<sup>1</sup> Kiel University

<sup>2</sup> German Aerospace Center (DLR)

#### References

- (1) P. Gege, "The water color simulator WASI: an integrating software tool for analysis and simulation of optical in situ spectra", Computers & Geosciences, Volume 30, Issue 5, June 2004, Pages 523-532
- (2) U. Müller-Wilm (2016), S2PAD SEN2COR 2.2.0 – Readme, S2PAD-VEGA-SRN-0001
- (3) ATCOR, Atmospheric/Topographic Correction for Airborne Imagery, ReSe Applications Schläpfer, Wil, Switzerland, 2011. [Online]. Available: www.rese-apps.com
- (4) H. Fientje, B. Briel, C. Beck, M. Collaud Coen, M. Fricke, J. Cyrys, J. Gu, M. Pittz, W. Thomas, "Identification and monitoring of Saharan dust: An inventory representative for south Germany since 1997", Atm. Env. 109, 87-96, 2015

#### Acknowledgements

This work was conducted within the project Kelpmap 2.0 (grant no: 50EE1020) funded by the Federal Ministry for Economic Affairs and Energy. We further thank ESA for providing Sentinel-2A data. We are very grateful to Katja Dornhöfer for helping out with toolboxes and templates.

Supported by:



on the basis of a decision by the German Bundestag



Contact:  
M. Sc. Sebastian Riedel  
EOM Research Assistant  
+49 (0) 8153 28 1959  
riedel@geographie.uni-kiel.de

Earth Observation and Modelling  
Kiel University  
Department of Geography  
Ludwig-Meyn-Str. 14  
24098 Kiel  
Germany